

## **Winter Manure Application Log**

Date of Application	Farm and Field #	Liquid or Solid	Spreader Capacity (gallons/tons)	<b>Loads</b> (# of trips)	Total Volume Applied (gallons/tons)	Acres Applied	Practices in Place When Required (select a practice letter below)

Slopes >6% select two practices: A-field has contour buffer strips or is in contour strip crop; **B**-all residue is left on the field and no fall tillage (does not apply where residue is removed as silage or bedding); **C**-apply on intermittent strips on no more than 50% of the field; **D**-apply to no more than 25% of the field waiting at least 14 days between applications; **E**-reduce application rates to 3,500 gal/ac or 30 lbs. P2O5 whichever is less.

**Concentrated Flow Channels select two practices**: Any of A-E above; **F**-no manure within 200 ft. of concentrated flow channels; **G**-fall tillage is on the contour and slopes are <6%.



## **Winter Manure Application Log**

Date of Application	Farm and Field #	Liquid or Solid	Spreader Capacity (gallons/tons)	<b>Loads</b> (# of trips)	Total Volume Applied (gallons/tons)	Acres Applied	Practices in Place When Required (select a practice letter below)
					_		

Slopes >6% select two practices: A-field has contour buffer strips or is in contour strip crop; **B**-all residue is left on the field and no fall tillage (does not apply where residue is removed as silage or bedding); **C**-apply on intermittent strips on no more than 50% of the field; **D**-apply to no more than 25% of the field waiting at least 14 days between applications; **E**-reduce application rates to 3,500 gal/ac or 30 lbs. P2O5 whichever is less.

**Concentrated Flow Channels select two practices**: Any of A-E above; **F**-no manure within 200 ft. of concentrated flow channels; **G**-fall tillage is on the contour and slopes are <6%.